

Algorithm for Gradual Hair Extensions, Taking into Account the Individual Characteristics of the Client, the Attachment Area, and Subsequent Care

Yuliia Cowan^{1*} 
¹Independent Researcher, Lviv, Ukraine

DOI: <https://doi.org/10.36348/sijcm.2026.v09i10.001>

| Received: 14.08.2026 | Accepted: 07.10.2026 | Published: 10.10.2026

*Corresponding author: Yuliia Cowan
Independent Researcher, Lviv, Ukraine

Abstract

Hair extension practice rarely fails in a single dramatic way. It fails gradually when the cumulative load on natural hair exceeds its structural reserve, when an attachment zone absorbs repeated tension it was not assessed to sustain, or when a client's actual maintenance behavior falls short of what the installation required. This article develops a literature-based algorithm for gradual hair extensions that treats the procedure as a staged adaptation process rather than a one-time intervention. The study draws on 20 verified English-language sources covering hair and scalp diagnostics, hair camouflage, scalp reactions, traction-related damage, and material exposure. It also references the Hair Biomechanical Tolerance Dataset (Cowan Y, 2026), a synthetic illustrative dataset of 108 hypothetical archetypes with rule-generated action directives. Three patterns from this model informed the algorithm. First, scalp condition was encoded as an absolute procedural gate: all archetypes with active erythema or pre-traction markers were assigned refusal, regardless of hair thickness, density, or zone. Second, staged extension is exclusively a Supportive and Conditional zone procedure: none of the 24 Staged Extension profiles is located in the high-risk marginal zone. Third, aftercare compliance reduces safe wear duration by 1.3 to 1.7 weeks across all scalp zones, making it a planning variable rather than post-procedural advice. On this basis, the article proposes a Gradual Hair Extension Algorithm with five decision blocks: structural diagnosis, zonal tolerance assessment, aftercare capacity assessment, phased load planning, and follow-up correction logic. Graduality is not a stylistic preference. It is a principle of biological proportionality.

Keywords: Hair extensions, gradual hair extensions, attachment area, aftercare, traction risk, hair cosmetics, scalp tolerance, beauty practice.

Copyright © 2026 The Author(s): This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC BY-NC 4.0) which permits unrestricted use, distribution, and reproduction in any medium for non-commercial use provided the original author and source are credited.

INTRODUCTION

Hair extension work is usually framed as a beauty service, yet its success depends on variables more typical of clinical or material assessment than of simple styling. A transformation may look convincing on the day of installation and still prove structurally excessive later, when the native hair must support added weight, the attachment areas must tolerate repeated movement, and the client must maintain the result through washing, brushing, drying, and correction visits. For that reason, the design of extensions cannot be evaluated solely by immediate appearance. It should also be evaluated by proportionality, durability, and the absence of avoidable biological cost.

This problem has become more relevant for two reasons. First, clients increasingly ask for dramatic change in a single appointment. They want more length, more fullness, more invisibility of fixation points, and a

result close to social-media references that often ignore differences in hair density, damage history, or scalp sensitivity. Second, the scientific and clinical literature surrounding hair, scalp, and cosmetic correction has grown considerably. There is now substantial knowledge about diagnostic evaluation of hair loss, cosmetic management of vulnerable hair, camouflage practices, contact reactions associated with hair products, synthetic-fiber exposure, and traction-related thinning (Dias M *et al.*, 2021; Singh S *et al.*, 2021; Daruwalla S *et al.*, 2022; Hwang J *et al.*, 2024; Abdallah S *et al.*, 2025). What remains underdeveloped is the translation of that knowledge into an applied extension algorithm.

The key issue is that extension practice is rarely governed by one variable alone. The individual characteristics of the client influence mechanical tolerance. The attachment area changes the local burden of the same donor mass. Subsequent care determines whether the installation remains orderly, dry, low-

friction, and safe between corrections. If one of these elements is underestimated, the procedure may be aesthetically impressive but functionally unstable. This is especially evident in clients with chemically treated hair, reduced density at the margins, scalp sensitivity, or limited ability to maintain a demanding care routine.

For that reason, this article proceeds from a different premise. It does not ask which extension method is best in general. It asks how a gradual extension strategy can be developed so that the degree of transformation is aligned with the carrying capacity of the client's hair, the tolerance of the attachment zone, and the reality of post-procedural care. The study draws on twenty verified English-language sources and on the Hair Biomechanical Tolerance Dataset (Cowan Y, 2026), which provides computed clinical action directives for 108 archetypes across the full range of structural profiles, scalp zones, and aftercare conditions. The aim is to formulate an evidence-informed algorithm suitable for professional consultation and staged extension planning.

The literature relevant to this topic is broad but fragmented. It includes work on hair and scalp diagnostics, aesthetic camouflage, prosthetic alternatives, scalp reactions to cosmetic products, synthetic-hair exposure, traction-related pathology, and texture-specific vulnerability. Individually, these publications answer different questions. Read together, they provide a useful foundation for a staged extension model.

The first group of studies concerns the assessment of hair and scalp status. Singh S et al (2021) propose a practical diagnostic approach to hair loss that emphasizes structured evaluation rather than impressionistic judgment. Ramsauer K (2021) likewise argues that management of hair concerns beyond standard therapy requires individualized reasoning. Kinoshita-Ise M et al (2022), writing on trichoscopy, reinforce the value of disciplined observation and shared terminology. Although these papers do not focus on extensions, they establish a principle essential for extension work: visible fullness is not the same as structural reserve. A consultation that ignores fragility, thinning, breakage, or scalp discomfort begins with incomplete information.

A second body of literature examines aesthetic compensation itself. Park J et al (2018) discuss wigs as a meaningful option in severe alopecia areata. Daruwalla S et al (2022) review hair camouflage in a broad practical sense, and Taye M et al (2023) consider camouflage in Black women while explicitly noting that some appearance-improving practices may worsen hair loss when they rely on adhesives, chemicals, or high-tension styling. Courtney A et al (2024) further show that wigs involve not only aesthetic but also psychosocial and economic dimensions. Together, these studies widen the

field of acceptable professional response. They imply that the desired look and the biologically safest method do not always coincide, and that a responsible specialist sometimes achieves the client's goal through reduction, redistribution, or substitution rather than through maximal durable attachment.

The third group of sources concerns reactivity, contact exposure, and the chemical environment of hair work. Karim M et al (2023) show that popular hair care practices and ingredients can create a meaningful potential for allergic contact dermatitis. Warshaw E et al (2022) summarize contact dermatitis associated with hair care products, Hwang J et al (2024) describe scalp allergic contact dermatitis as an underdiagnosed problem, and Cebolla-Verdugo M et al (2024) review contact dermatitis due to hair care products more broadly. These papers matter because the extension environment is not defined only by fixation. It is also defined by intensified product exposure, longer drying times, friction patterns around bonds or rows, and sometimes adhesives or synthetic fibers close to the scalp. The literature on synthetic materials strengthens this point. Dlova N et al (2021) report irritant contact dermatitis associated with synthetic hair extensions in patients with an atopic history. Auguste D et al (2020) show volatile organic compound emissions from heated synthetic hair, and Franklin E et al (2026) identify chemicals of health concern in extension materials. Such findings do not invalidate hair extensions as a category, but they show that material choice and heat behavior belong inside the risk assessment.

The final major literature cluster concerns traction and region-specific vulnerability. Abdallah S et al (2025) report a high prevalence of traction alopecia in women in North Sudan and identify associated factors connected with hair practices. Anaba E et al (2025) demonstrate measurable reductions in hair density and shaft diameter in different severities of anterior scalp marginal traction alopecia. Geisler A et al (2024), reviewing disorders in curly hair, emphasize that texture-specific vulnerability, grooming patterns, and mechanical stress can alter both diagnosis and management. Chen L et al (2024), although presenting a case-based discussion, also remind us that altered texture behavior may signal a bigger change. Taken together, these sources suggest that the margin and other sensitive attachment areas must be treated as zones of limited tolerance, especially when density is already reduced or the hair is fragile, textured, or cosmetically compromised.

One more important issue appears in research on client perceptions of corrective options. Kim M et al (2025), studying scalp micropigmentation among dermatology outpatients with hair loss, show that awareness, preference, and treatment experience shape how clients interpret aesthetic correction. This is directly relevant to consultation in extension practice. Clients do

not necessarily request the most appropriate solution; they request the form of correction they know. A professional algorithm, therefore, has to translate visual desire into a safer procedural equivalent.

An illustrative quantitative example of this gap is available. The Hair Biomechanical Tolerance Dataset (Cowan Y, 2026) computes a hypothetical clinical action directive for each of 108 client archetypes using four integrated factors: base carrying capacity derived from shaft diameter, a degradation coefficient reflecting chemical history, anatomical tension mapping across scalp zones, and clinical modifiers including scalp condition and aftercare compliance. The dataset's three-tier zonal classification, Supportive, Conditional, and High-Risk, directly parallels the zone-based logic that the traction literature implies. Its Clinical_Action outputs, which range from Standard Single-Stage through Staged Extension to Refusal, make explicit what the reviewed literature implies: that the appropriate procedural response is not uniform across clients, and that the transition between procedural categories can be sharp rather than gradual.

The central gap is now clear. There is substantial knowledge about vulnerability, camouflage, materials, and correction, but there is still limited integrated guidance for deciding when enhancement should be immediate, when it should be phased, and when it should be redirected to a lower-burden option. That gap justifies the present attempt to build an extension algorithm grounded not in method prestige but in compatibility among the client profile, the attachment area, and the subsequent care model.

MATERIALS AND METHODS

This study is conceptual and analytical, but it includes a small secondary empirical component. No

primary salon testing, no interviews, and no laboratory measurements were conducted for this article. The evidence base consisted of 20 verified English-language sources selected for direct relevance to at least one of four domains: diagnostic evaluation of hair and scalp, aesthetic correction or camouflage, material and product-related reactivity, and traction or texture-specific vulnerability.

The analytical procedure had three stages. First, the literature was classified by problem type. Diagnostic papers clarified how clinicians and specialists distinguish visible fullness from actual structural reserve (Singh S *et al.*,2021; Ramsauer K, 2021; Kinoshita-Ise M *et al.*,2022). Camouflage and prosthetic papers clarified that not all aesthetic correction needs to be load-bearing or fixed (Park J *et al.*,2018; Daruwalla S *et al.*,2022; Taye M *et al.*,2023; Courtney A *et al.*,2024). Reactivity studies identified risks associated with hair products, synthetic fibers, and scalp exposure (Karim M *et al.*,2023; Hwang J *et al.*,2024; Cebolla-Verdugo M *et al.*,2024; Dlova N *et al.*,2021; Auguste D *et al.*,2020; Franklin E *et al.*,2026). Traction- and texture-related papers clarified which zones and hair profiles are more vulnerable to load and manipulation (Abdallah S *et al.*,2025; Anaba E *et al.*,2025; Geisler A *et al.*,2024; Chen L *et al.*,2024).

Second, these studies were reinterpreted through the logic of extension consultation. The question was not what each article originally set out to prove, but which repeated variables matter most when planning safer enhancement. Six variables were retained: structural reserve of the natural hair, density reserve, texture behavior, attachment-area tolerance, material or scalp reactivity, and expected aftercare compliance. These were then converted into practical consultation dimensions.




Figure 1: Decision logic of the Gradual Hair Extension Algorithm, including the scalp condition gate and zone-based restrictions on staged extension

Third, the study integrated the dataset (Cowan Y, 2026) as an empirical reference for the algorithm's decision boundaries. The dataset archetypes were examined to verify five propositions that the literature supports qualitatively: that compromised hair or scalp status lowers tolerance to heavy enhancement; that marginal and tension-prone zones carry increased risk; that material choice affects comfort and safety; that some clients are better served by lower-burden corrective options; and that maintenance behavior modifies outcome stability. The computed outputs of the illustrative dataset were consistent, by construction, with all five propositions drawn from the literature, and the pattern of results, particularly the zone-based and scalp-condition-based boundaries between procedural categories, directly shaped the structure of the proposed algorithm. It should be noted that the referenced dataset

consists of illustrative synthetic archetypes generated to explore parameter relationships, and does not represent empirical observations from real clients or patients.

The final methodological step was model construction. Instead of mapping the algorithm to commercial brands or to one proprietary extension system, the article describes extension options through behavior categories: higher-burden versus lower-burden correction, broader versus more localized distribution of load, reversible versus prolonged wear, and immediate versus phased transformation. This makes the model transferable across salon contexts. It also aligns better with the literature, which speaks more consistently about vulnerability and tolerance than about brand-specific methods.

Table 1: Analytical structure of the gradual extension methodology

Analytical block	Core variables	Methodological function
Client profile	Density, thickness, fragility, cosmetic history, scalp sensitivity	Defines the biological carrying capacity of the natural base.
Attachment area	Marginal zones, mobile zones, supportive zones, visibility requirements	Determines where durable or reduced-load attachment is acceptable.
Care assessment	Brushing discipline, hygiene routine, heat styling habits, and correction compliance	Predicts whether the client can maintain the planned enhancement safely over time.
Dataset reference layer	Hair Biomechanical Tolerance Dataset (Cowan Y, 2026), 108 archetypes, computed clinical action directives	Identifies precise conditions under which Staged Extension, High-Risk Protocol, or Refusal is indicated; confirms zone-based and scalp-condition-based boundaries

RESULTS AND DISCUSSION

The synthesis produced three principal findings. First, safe extension planning depends less on selecting a fashionable method than on matching the scale of enhancement to the carrying capacity of the natural hair. Second, the attachment area is a decisive predictive variable rather than a minor technical detail. Third, aftercare capacity should be treated as part of the procedural decision itself.

The literature supports a basic classification of clients by tolerance profile. One profile is relatively resilient: acceptable density, limited visible fragility, no major scalp sensitivity, and a care routine likely to be maintained reliably. A second profile is conditional: the hair can support enhancement, but only if the burden is reduced and zonal placement is selective. A third profile is vulnerable: low density, edge weakness, strong chemical history, curly-hair fragility under manipulation, scalp reactivity, or poor expected compliance with aftercare. In this third profile, the literature repeatedly favors lower-burden correction, staged enhancement, or alternative camouflage rather than ambitious immediate installation (Daruwalla S *et al.*, 2022; Taye M *et al.*, 2023; Geisler A *et al.*, 2024).

Dataset analysis reinforced and quantified this picture. The most decisive finding concerns scalp condition as a procedural gate. Every archetype in the dataset with active erythema or pre-traction markers received a Refusal directive, independent of hair thickness, chemical history, or any other variable, six erythema profiles and five pre-traction profiles, all resulting in 100% refusal. By contrast, only 8% of profiles with a healthy scalp condition were refused. This finding provides computational confirmation for what the traction literature implies but rarely states numerically: scalp condition is not one variable among many. It is a gate that must be checked before any other assessment proceeds. The zone-based findings were equally clear. Staged Extension appeared in 44% of Supportive zone profiles and 21% of Conditional zone profiles, making it the most common directive in the former. In the High-Risk zone, Staged Extension appeared in 0% of profiles. The High-Risk zone distributed its cases among Careful Monitoring (47%), High-Risk Protocol (19%), and Refusal (33%), with Standard Single-Stage also absent. The algorithm's zonal logic, which restricts staging to Supportive and Conditional zones, is directly supported by this pattern (Figure 2).

Clinical action distribution by scalp zone




Figure 2: Clinical action distribution by scalp zone (n = 108 archetypes). Staged Extension is absent from the High-Risk zone

Aftercare compliance operated differently depending on the zone and scalp condition. It did not shift profiles between procedural categories in a systematic way, but it consistently shortened the safe wear cycle within each category. In the Supportive zone, the mean wear cycle fell from 7.8 weeks with high compliance to 6.1 weeks with low compliance, a reduction of 1.7 weeks. In the Conditional zone, the gap

was 1.6 weeks (6.7 vs 5.1 weeks). Even in the High-Risk zone, where wear cycles were already the shortest, low compliance removed a further 1.3 weeks relative to high-compliance profiles (5.7 vs 4.4 weeks, Figure 3). These data support treating aftercare compliance as a temporal planning variable: it does not determine which procedural category a client enters, but it determines how quickly that procedure must be followed by correction.

Wear cycle by scalp zone and aftercare compliance




Figure 3: Mean wear cycle by scalp zone and aftercare compliance. Refusal rates for each zone are shown at the base of the chart

On this basis, the authorial Gradual Hair Extension Algorithm was developed (Figure 1). The first block is structural diagnosis. The specialist evaluates density, thickness, visible fragility, cosmetic history, texture behavior, and signs of scalp discomfort. The second block is zonal assessment. The scalp is classified into more supportive, conditionally supportive, and high-risk areas. The third block is aftercare assessment. The practitioner evaluates whether the client can realistically maintain washing, drying, separation of attachment points, low-friction styling, and timely correction appointments. The fourth block is phased load planning. Instead of moving directly to the maximum desired transformation, the specialist decides whether the case calls for a conservative single-stage installation, a two-stage progressive increase, a multi-stage adaptation plan, or redirection toward a removable or lower-burden solution. The fifth block is follow-up correction logic. Reassessment is built into the method before any increase in mass, length, or density is approved.

The distinctive feature of the algorithm is that it treats graduality as a professionally precise response, not a compromise forced by indecision. Dataset analysis quantifies where this precision matters most. In the High-Risk zone, a third of profiles result in outright refusal. In the Supportive zone, nearly half can undergo staged extension safely, meaning the same principle of caution

produces fundamentally different plans depending on anatomical placement. The algorithm captures this by making zone assessment a mandatory second block: the plan cannot advance to load decisions until the attachment area has been classified. If a client requests prominent perimeter framing in a marginal zone already showing edge weakness, the algorithm does not simply reduce the planned mass. It redirects the visual task to a zone that can support it or recommends a removable alternative for the marginal area.

Aftercare capacity is the third pillar of the algorithm, and the dataset confirms its planning role in concrete terms. A difference of 1.3 to 1.7 weeks in safe wear duration between high- and low-compliance profiles means that two clients with identical hair and the same attachment zone may need appointments scheduled at different intervals. For a Supportive-zone client with compromised hair and low compliance, the computed wear cycle drops to approximately four to five weeks, the minimum correction interval in the dataset. At that point, the algorithm must either confirm that the client can realistically meet this schedule or reduce the installation scope until the required correction frequency becomes manageable. Thus, the aftercare assessment is not a conversation that follows the procedure. It is the reasoning that determines whether the procedure should proceed at its planned scale at all.

Table 2: Decision blocks of the Gradual Hair Extension Algorithm and their practical effects

Algorithmic block	Core action	Expected practical effect
1. Diagnosis	Assess density, shaft thickness, fragility, cosmetic history, and scalp tolerance.	Establishes whether the client can safely support enhancement.
2. Zone mapping	Classify attachment areas as supportive, conditional, or high-risk.	Reduces local overload and marginal tension.
3. Care screening	Evaluate whether the client can maintain brushing, hygiene, heat control, and timely correction.	Prevents a technically correct installation from failing during wear.
4. Gradual load choice	Choose one-stage conservative enhancement or a staged progression.	Aligns aesthetic ambition with structural tolerance.
5. Follow-up schedule	Plan correction timing and observe comfort, shedding, and scalp reaction.	Turns the method into a monitored long-term strategy rather than a one-time procedure.

The proposed model reframes hair extensions as a longitudinal compatibility problem. This is one of its main theoretical and practical contributions. In common salon discourse, success is often measured by immediate visual transformation. The literature reviewed here suggests that such a standard is too narrow. An extension result should also be judged by whether it remains proportionate across wear, care, and correction cycles. A method that looks seamless on day one but overloads vulnerable areas or exceeds the client's maintenance capacity cannot be considered fully successful.

The article also clarifies why graduality is scientifically meaningful. It is not simply a softer aesthetic philosophy. It is a way of reducing uncertainty in a context where the variables are interdependent. The structure of the native hair, the chosen attachment area,

and the subsequent care environment all modify each other. A phased strategy allows the practitioner to test tolerance before authorizing greater change. In that sense, graduality functions much like dose adjustment in other applied fields: one begins proportionately, observes response, and only then increases burden where justified.

The reviewed literature on camouflage and prosthetic options is especially helpful here. Park J et al (2018), Daruwalla S et al (2022), Taye M et al (2023), and Courtney A et al (2024) all imply that aesthetic correction exists on a continuum. Once this is accepted, the specialist is no longer forced into a binary choice between "full extensions" and "no correction." The field of professional response becomes wider and more honest. This is important because some of the greatest

technical errors in beauty practice occur when the practitioner tries to satisfy an aesthetic request with a procedure that the substrate cannot safely support.

Attachment-area logic deserves the same emphasis. Traction-related literature shows that not all scalp zones are equally forgiving (Abdallah S *et al.*, 2025; Anaba E *et al.*, 2025). In practical terms, this means that uniform installation maps are difficult to defend scientifically. If a margin is already fragile, the correct response is not to hide the problem beneath more ambitious fixation. It is to protect the zone, reduce local demand, or move the visual task elsewhere. The same principle applies to clients with curly textured hair or altered texture behavior, where grooming stress and fragility may already be elevated (Geisler A *et al.*, 2024; Chen L *et al.*, 2024).

Aftercare is the third pillar of the model. Product-related and scalp-reaction studies strongly suggest that maintenance behavior changes the risk environment of any enhancement procedure (Karim M *et al.*, 2023; Warshaw E *et al.*, 2022; Hwang J *et al.*, 2024; Cebolla-Verdugo M *et al.*, 2024). If a client is unlikely to dry attachment areas carefully, avoid excessive heat, return on time for correction, or maintain low-friction handling, then even a well-chosen installation may become unsafe. The practical implication is clear: a realistic care conversation belongs before the procedure, not after it.

The dataset confirms this logic numerically. Aftercare compliance reduced mean wear cycles by 1.3 to 1.7 weeks across all zones, independently of hair structure or chemical history. In a zone where the starting wear cycle is already short, such as the High-Risk zone at 5.7 weeks for high-compliance profiles, a further reduction to 4.4 weeks for low-compliance clients means that the correction window compresses to the absolute minimum. At that point, the realistic option is not a better installation technique but a frank conversation about whether the planned procedure remains compatible with how this client actually lives with their hair.

The article has limitations. It does not include a prospective salon trial, and the dataset archetypes are computationally modeled rather than drawn from live client observations. They provide reliable structural reference points, but they do not capture individual variation in pain thresholds, actual home care behavior, or scalp reactivity to specific product formulations. The proposed algorithm should therefore be understood as an evidence-informed consultation framework rather than a predictive clinical instrument. Another limitation is that some of the most useful source publications are adjacent to extensions rather than extension-specific. Yet this reflects the reality of the field: real extension practice is shaped simultaneously by traction diagnostics, reactivity, camouflage logic, and client behavior, and a

narrowly technical literature review would not capture that complexity.

Future research should test the algorithm in salon settings through observational or prospective designs. It would be especially useful to compare immediate full-load enhancement with phased enhancement across different tolerance profiles and to examine whether exclusion of high-risk attachment zones improves long-term outcomes. Simplified scoring systems for structural reserve, zonal tolerance, and aftercare reliability could also make the model easier to standardize in education and practice.

CONCLUSION

The analysis supports the conclusion that gradual hair extensions should be treated as an adaptive and risk-sensitive strategy rather than as a slower version of ordinary installation. Their scientific value lies in proportionality. Individual characteristics of the client, attachment-area tolerance, and subsequent care capacity operate together as a single decision system. When one of these dimensions is ignored, the visible result may still be attractive, but the procedure becomes less stable and potentially more harmful over time.

The main contribution of this article is the formulation of the Gradual Hair Extension Algorithm derived from the synthesis of 20 verified English-language sources and grounded in the Dataset (Cowan Y, 2026). The model organizes extension planning into five linked blocks: structural diagnosis, zonal assessment, aftercare assessment, phased load planning, and follow-up correction logic. Analysis of the illustrative dataset demonstrates three decision boundaries that the algorithm formalizes, reflecting the model's built-in assumptions rather than independent clinical confirmation: erythema and pre-traction markers produced 100% refusal regardless of all other parameters. Staged Extension is exclusively a Supportive and Conditional zone procedure: not one high-risk zone profile in the dataset received a staged directive. Aftercare compliance reduces safe wear cycle duration by 1.3 to 1.7 weeks, making it a planning variable that belongs in consultation, not in post-installation advice. Together, these boundaries define the conditions under which graduality is appropriate, where it must give way to more conservative options, and where the procedure cannot proceed at any scale.

In practical terms, this means that the most professional extension strategy is not necessarily the most dramatic one. It is the one that creates a convincing appearance while preserving the client's own hair and scalp over time. In theoretical terms, the article helps integrate dermatological, cosmetic, and aesthetic evidence into a single framework directly relevant to extension practice. Future validation is still required, but the logic of the model is already clear: in advanced

beauty work, responsible transformation is gradual because the hair must first be able to carry it.

REFERENCES

- Dias, M. F. R. G., Loures, A. F., Ekelem, C., *et al.*, (2021). Hair cosmetics for the hair loss patient. *Indian Journal of Plastic Surgery*, 54(4), 507-513. <https://doi.org/10.1055/s-0041-1739241>
- Singh, S., Kumar, M., Bala, M., & Khare, A. K. (2021). Practical approach to hair loss diagnosis. *Indian Journal of Plastic Surgery*, 54(4), 514-521. PMID: 34984076.
- Daruwalla, S. B., Dhurat, R. S., & Hamid, S. A. T. (2022). All that a dermatotrichologist needs to know about hair camouflage: A comprehensive review. *International Journal of Trichology*, 14(3), 77-83. https://doi.org/10.4103/ijt.ijt_6_21
- Hwang, J. C., Beatty, C. J., Khobzei, K., *et al.*, (2024). Allergic contact dermatitis of the scalp: A review of an underdiagnosed entity. *International Journal of Women's Dermatology*, 10(3), e167. PMID: 39076889.
- Abdallah, S., *et al.*, (2025). Prevalence and associated factors of traction alopecia in women in North Sudan: A community-based cross-sectional study. *International Journal of Women's Dermatology*. PMID: 40005312.
- Cowan, Y. (2026). Research Toolkit Biomechanical Tolerance Dataset and Computational Model for Safe Hair Extension Architecture (1.0) [Data set]. Zenodo. <https://doi.org/10.5281/zenodo.21037485>
- Ramsauer, K. C. (2021). A practical guide to treatment of hair loss beyond standard therapy. *Cutis*, 107(4), E11-E15. <https://doi.org/10.12788/cutis.0236>
- Kinoshita-Ise, M., *et al.*, (2022). Update on trichoscopy: Integration of the terminology by systematic review of the literature and consensus of the International Trichoscopy Society. *Journal of the European Academy of Dermatology and Venereology*. PMID: 34806223.
- Park, J., Kim, D.-W., Park, S.-K., Yun, S.-K., & Kim, H.-U. (2018). Role of hair prostheses (wigs) in patients with severe alopecia areata. *Annals of Dermatology*, 30(4), 505-507. <https://doi.org/10.5021/ad.2018.30.4.505>
- Taye, M., Kamili, N. A., Aguh, C., & Krueger, L. D. (2023). Considerations and recommendations on camouflage in alopecia in Black women. *International Journal of Women's Dermatology*, 9(3), e111. <https://doi.org/10.1097/JW9.0000000000000111>
- Courtney, A., & Su, J. C. (2024). Wigs and alopecia areata: Psychosocial impact and economic considerations. *Cosmetics*, 11(2), 55. <https://doi.org/10.3390/cosmetics11020055>
- Karim, M., Klein, E. J., Nohria, A., *et al.*, (2023). Potential for allergic contact dermatitis in popular hair care practices and ingredients. *Dermatitis*, 34(6), 484-491. PMID: 37339433.
- Warshaw, E. M., *et al.*, (2022). Contact dermatitis associated with hair care products. *Dermatitis*. PMID: 34149001.
- Cebolla-Verdugo, M., Velasco-Amador, J. P., & Navarro-Triviño, F. J. (2024). Contact dermatitis due to hair care products: A comprehensive review. *Cosmetics*, 11(3), 78. <https://doi.org/10.3390/cosmetics11030078>
- Dlova, N. C., Ferguson, N. N., Rorex, J. N., *et al.*, (2021). Synthetic hair extensions causing irritant contact dermatitis in patients with a history of atopy: A report of 10 cases. *Contact Dermatitis*, 84(5), 357-359. <https://doi.org/10.1111/cod.13825>
- Auguste, D., *et al.*, (2020). Volatile organic compound emissions from heated synthetic hair: A pilot study. *Environmental Health Insights*, 14. PMID: 32063708.
- Franklin, E. T., Favela, K., Spies, R., Ranger, J. M., & Rudel, R. A. (2026). Identifying chemicals of health concern in hair extensions using suspect screening and nontargeted analysis. *Environment & Health*. <https://doi.org/10.1021/envhealth.5c00549>
- Anaba, E. L., & Sani, H. (2025). A comparative cross-sectional study of hair density and hair shaft diameter of different severities of anterior scalp marginal traction alopecia in women. *Skin Appendage Disorders*. PMID: 40475102.
- Geisler, A. N., Taye, M., Larrondo, J., Mayo, T. T., Aguh, C., McMichael, A., MacKelfresh, J. B., & Krueger, L. (2024). Updates on disorders in curly hair. *International Journal of Dermatology*, 63(9), 1145-1154. <https://doi.org/10.1111/ijd.17184>
- Chen, L.-C., & Aguh, C. (2024). "Downy hair sign"-A clue to autoimmune disease in curly hair. *JAAD Case Reports*, 51, 59-62. <https://doi.org/10.1016/j.jdc.2024.06.018>
- Kim, M. J., & Kim, H. S. (2025). Perceptions and experiences of scalp micropigmentation among dermatology outpatients with hair loss: A survey-based study. *Annals of Dermatology*. PMID: 41331716.