

Polycarbonate Bracket with Herbal Dentifrices – Comparative Clinical and Microbiological Study

Dr. Aseem Sharma¹, Dr. Sachin Upadhyay^{2*}, Dr. Gaurav Didhra³, Dr. Mandeep Nain⁴, Dr. Yatharth Goel⁵, Dr. Neelam Chauhan⁶

¹Senior Lecturer in Department of Orthodontics and Dentofacial Orthopedics at Himachal Institute of Dental Sciences Paonta Sahib, Himachal Pradesh India

²Senior Lecturer in Department of Orthodontics and Dentofacial Orthopedics at Himachal Institute of Dental Sciences Paonta Sahib, Himachal Pradesh India

³Director at Dent-O-Care, Opp UCO Bank Main bazaar, Mehatpur, Una India

⁴Post Graduate Student in the Department Of Prosthodontics, DAV Dental College, Yamuna Nagar

⁵Director at Dantam Dental Solutions 44, civil lines, Jadugar Road, Roorkee India

⁶Consultant Orthodontist at Kullu Smiles, Kullu Himachal Pradesh India

DOI: 10.36348/SJODR.2019.v04i09.012

| Received: 11.09.2019 | Accepted: 20.09.2019 | Published: 30.09.2019

*Corresponding author: Dr. Sachin Upadhyay

Abstract

Background: To ascertain if herbal dentifrices have an effect on Streptococcus mutans count in orthodontic patients with Polycarbonate brackets. **Material and method:** patient had tooth No's 45 included in the study with polycarbonate bracket bonded. Dentifrices tested were Herbal based. Conventional tooth paste was considered as control group. **Result:** Paired T test compared the means of Streptococcus mutans count around polycarbonate bracket at different time intervals. **Conclusion:** This shows polycarbonate bracket has statistically significant reduction of Streptococcus mutans with herbal toothpaste.

Keywords: Polycarbonate Bracket, Herbal, Streptococcus Mutans.

Copyright © 2019: This is an open-access article distributed under the terms of the Creative Commons Attribution license which permits unrestricted use, distribution, and reproduction in any medium for non-commercial use (NonCommercial, or CC-BY-NC) provided the original author and source are credited.

INTRODUCTION

Unlike the majority of oral microorganisms, Streptococcus mutans thrives under acidic conditions and becomes the dominant bacterium in cultures with permanently reduced pH [2]. Though, brushing teeth twice a day is considered reasonably effective in plaque and bacterial count reduction, the common prevalence of gingival inflammation in orthodontic patients often suggests inadequate oral hygiene procedures in most patients. Meswak *Salvadorapersica* (Meswak) is a medicinal herbal plant which has been used for centuries as oral hygiene tools. Chemical analysis of *S. persica* demonstrated the presence of many components exhibiting antimicrobial effect. Meswak (*Salvadorapersica*) contains salvadorine and trimethylamine, which are shown to exhibit antibacterial effects on cariogenic [3] bacteria such as *Streptococcus mutans*. It has been shown that these active principles support periodontal health, reduces the accumulation of biofilm-like dental plaque formation and exhibits fungistatic activity against *Candida albicans*[3]. Meswak is available in two forms namely meswak sticks and meswak tooth paste. Pomegranate is currently finding important applications in the field of

dental health. Clinical studies [5] have shown that this popular antioxidant attacks the causes of tooth decay at the biochemical level, with remarkable vigour. The fruit of the pomegranate tree has been used extensively in the folk medicine of many cultures. The healing property of pomegranates was discussed in one of the oldest medical texts, the Ebers Papyrus from ancient Egypt (1500 BC) [5]. Hence, this study of microbiological assay of Streptococcus mutans with an objective to study the performance and measure the efficacy of toothpastes- Neem, Meswak, Babool and Pomegranate based herbal toothpaste with polycarbonate Bracket is done.

AIM AND OBJECTIVES

To ascertain if herbal dentifrices have an effect on Streptococcus mutans count in orthodontic patients with Polycarbonate brackets.

MATERIAL AND METHOD

Nature of Study

Randomized, prospective, cross sectional single blinded microbiological assay study with each patient acting his /her own control in this study.

Area of Study

Department of Orthodontics and Dentofacial Orthopedics, Divya Jyoti College of Dental Sciences and Research and Microbiological Assay were conducted in Divya Jyoti Hospital.

Ethical Clearance

This study was approved by Institutional Committee (IEC No DJD/IEC/2014/A-001). A written consent was taken from each participating subject.

Inclusion criteria

- Patient with similar socioeconomic strata & common food habits.
- Patients free of oral/parental antibiotics for one month.
- No periodontal & systemic disorders
- Patients with no crowding and who have had alignment and leveling completed.

Dentifrices

S.No	Details	Code
1	Himalaya Herbals Dentifrice (Himalaya Global Holding Ltd.) Containing Neem, Meswak Babool and Pomegranate	Y (Blue)

Bracket Type

S. No	Bracket type
1	Polycarbonate Bracket [0.22 slot (Rhomboidal) MBT Libral Traders]



Polycarbonate Bracket

Steps and Time Interval of Study

- Each group consists of 30 teeth with 30 brackets to be tested.
- Each patient served as his/her own control as 1 types of bracket were tested in the same mouth at the same time period.
- Each patient had tooth No's 45 included in the study with polycarbonate bracket bonded.
- Dentifrices tested were Herbal based.
- The dentifrices were dispensed into 5ml bottles coded as Y for Herbal toothpaste –Neem, Babool, Meswak and Pomegranate. Color Coding of Dentifrices
- Conventional tooth paste was considered as control group.

S. No	N	Type	Bracket Bonded on Tooth Number
I	30	Polycarbonate Rhomboidal MBT	45



CODES	COLOR	DENTIFRICES
Y	Blue	Herbal Based Toothpaste (Herbal Global Holding Ltd Toothpaste)



Ice Box Plaque Collection and Transportation

Plaque collection and transportation

- Plaque sample placed in 5ml sterilized vials with 1ml distilled water.
- Sterilized vials were transported in icebox to the lab.
- The bacteriological study was conducted by Dilution Plating Method.
- The growth media used was Mutans-Sanguis Agar.

Oral Hygiene Instructions

- The subjects were given oral hygiene instructions & requested to refrain from using any other oral hygiene products like mouthwash etc.
- The subjects were instructed to follow standard oral hygiene regime which included brushing twice a day with toothpaste as prescribed in the study regime.
- The patients were advised to rinse thoroughly after every meal.

Table Shows: Time Interval of Tooth Paste Usage

TOOTHPASTE	TIME INTERVAL
Baseline without use of study Dentifrices	1 st to 2 nd Day
Herbal (Y)	3 rd to 8 th Day

**Table Shows: Time Interval of Plaque Collection
Plaque Collection Method**

Sample Count	Time Interval	Day Count
Sample No.1 (baseline without use of study dentifrices)	(T ₁) (Start of study)	Day : 1
Sample No.2	T ₂	Day : 3
Sample No.3	T ₃	Day : 8

- Patients were requested to refrain from eating or drinking 1 hour prior to sample collection.
- Plaque sample was collected by Four Pass Technique at midmorning (11 a.m.).
- In this technique the explorer tip is moved around the circumference of the bracket at the bracket tooth interface.
- Four passes, along the tooth at the bracket interface at the gingival, mesial, distal, and occlusal aspects are done to avoid overloading the instrument tip.
- This is considered an effective method of obtaining the total plaque. Plaque samples were placed in sterilized vials having distilled water in it.



Laboratory Equipment's



Mutans Sanguis Agar



Laminar Air Flow



Sterilization of Diluted Agar Medium in Autoclave



Distilled Water



Petridishes Placed Inside Incubator



Wire Loop




Solidification of Agar Medium in Laminar Air Flow



Micropipette



Incubator



Spreading of Plaque Sample over Petridish

S.No	Item
1	Autoclave
2	Hotplate
3	Petridish
4	Micropipette
5	Laminar flow Cabinet
6	Conical flask
7	Cotton Plug
8	Sterilized Wire loop
9	Incubator
10	Disposable gloves
11	U shape flask
12	Disposable Mouth mask

RESULT

Table Shows: Comparison of Means of Streptococcus mutans Count at Different Time Intervals around Polycarbonate Bracket by Paired T – Test.

Days			Mean difference	T	d.f.	P value
Day 1	-	Day 3	0.10000	0.619	29	0.541*
Day 3	-	Day 8	0.83333	5.473	29	0.000***

***Highly Significant $p < 0.001$, **Significant $p < 0.05$, *Not Significant $p > 0.05$

- Paired T test compared the means of Streptococcus mutans count around polycarbonate bracket at different time intervals.
- Difference between Day 3 & Day 8 with herbal dentifrice was highly significant statistically
- Difference between Day 1 & Day 3 was not statistically significant.
- The mean difference between Day 3 & Day 8 is 0.83333.

CONCLUSION

- This shows polycarbonate bracket has statistically significant reduction of Streptococcus mutans with herbal toothpaste.

DISCUSSION

The increased time and difficulty of maintaining good oral hygiene during orthodontic treatment are challenges faced by patients and the levels of oral bacteria have been reported to increase five folds due to the orthodontic devices and attachments [1]. *Emilson CG*. Stated that most orthodontic patients are not able to perform effective plaque control, and therefore develop mild to moderate gingivitis during treatment with fixed appliances. Microorganisms play a major role in causation of WSL and dental caries. Entire removal of microorganism from the oral cavity is difficult but their count can be reduced with the help of various preventive measures so that it becomes less cariogenic. The market is flooded with numerous bracket types of different biomaterials. Literature evidences that adherence of plaque to the fixed appliance is largely contributed by the bracket material [2] as it could play a role in the degree of bacterial adhesion and plaque accumulation as well as in the risk of development of WSL. The initial affinity of bacteria to solid surfaces is due mostly to electrostatic and hydrophobic interactions. Surfaces with high surface free energy more easily attract bacteria such as *S.mutans*[3]. Currently Brackets are being driven by manufacturers as having a hygiene advantage, while many studies have reported in the contrary [5]. The results of the current research study showed significant reduction around polycarbonate bracket with herbal toothpaste. The value of current study suggests that herbal dentifrices have good antimicrobial effects on caries producing bacteria, thus can be used in orthodontic patients and as a regular home care preventive aid in combating dental caries.

REFERENCES

- Papaioannou, W. (2007). Adhesion of Streptococcus mutans to Different Types of Brackets. *Angle Orthod*, 77(6) 1090-1095.
- Gastel, J.V., Quirynen, M., Teughels, W. (2009). Microbial Adhesion on Different Bracket Types In vitro study. *Angle Orthod*, 79:915–921.
- Brandão, G.A.M., Pereira, A.C., Brandão, A.M.M. (2015). Does the bracket composition material influence initial biofilm formation? *Indian J Dent Res*, 26:148-51.
- Rundell, B.B., Thomson, L.A., Stile, H.M. (1937). Evaluation of a new transport medium for the preservation of oral streptococci. *Archs oral Biol*, 18:871-878.
- Emilson, C.G. (1983). Prevalence of Streptococcus mutans with different colonial morphologies in human plaque and saliva. *Scand J Dent Res*, 91(1):26-32.

7. Tanzer, J.M., Borjesson, C., Laskowski, L., Testa, M. (1984). Glucose –Sucrose-Potassium Tellurite-Bacitracin Agar, an alternative to Mitis Salivarius –Bacitracin Agar for enumeration of *Streptococcus mutans*. J Clin Micro, 20(4): 653-659.
8. Mohammad, M. A. (2015). Review of the therapeutic effects of using Meswak (*Salvadora Persica*) on oral health Saudi Med J, 36(5): 530–543.
9. Sukontapatipark, W., Agroudi, M.A. (2001). Bacterial colonization associated with fixed orthodontic appliances. Eur J Orthod. 23(5):475-84.
10. Ahn, S.J., Kho, H.S., Lee, S.W. (2002). Roles of salivary proteins in the adherence of oral streptococci to various orthodontic brackets. J Dent Res. 81(6):411-415.
11. Kote, S., Nagesh, L. (2011). Effect of Pomegranate Juice on Dental Plaque Microorganisms (Streptococci and Lactobacilli). Anc Sci Life, 31(2): 49–51.
12. Citra, L., Yuwono, I., Benny, M. (2012). Effectiveness of herbal and non-herbal toothpastes in reducing dental plaque accumulation. J Dentistry Indonesia, 19: 32-40.
13. Halawany, H.S. (2012). A review on Meswak (*Salvadora persica*) and its effect on various aspects of oral health. Saudi Dent J, 24(2): 63–69.